

QUASI-REGULAR REPRESENTATIONS AND PROPERTY RD

ADRIEN BOYER

ABSTRACT. We give new formulations of property RD in terms of an L^1 -integrability condition of a Banach representation. This enables us to apply the theorem of Banach-Steinhaus and leads to a simple condition for property RD. We deduce that the quasi-regular representation on the Furstenberg boundary of any non compact semisimple Lie group with finite center has a matrix coefficient which does not verify the *weak inequality* of Harish-Chandra.

1. INTRODUCTION

The property of rapid decay has been introduced by H. Haagerup at the end of the seventies in his work [7]. The terminology “property RD” was introduced later, see the work [9] of P. Jolissaint. Property RD is relevant in the context of the Baum-Connes conjecture thanks to the important work of V. Lafforgue in [10].

We say that a unitary representation $\pi : G \rightarrow U(H)$ in a Hilbert space of a locally compact group with a length function L (see Definition 3.1), has property RD with respect to L if there exists $d \geq 1$ and $C > 0$, such that π verifies for all $f \in L^1(G)$,

$$\|\pi(f)\| \leq C \|f\|_{L,d}$$

where $\|\cdot\|$ denotes the operator norm, and $\|f\|_{L,d} = \sqrt{\int_G |f(g)|^2 (1 + L(g))^{2d} dg}$, where dg denotes a Haar measure on G . Equivalently, thanks to an observation from M. Perrone in [12], a representation $\pi : G \rightarrow U(H)$ has property RD with respect to L if there exists $d \geq 1$ and $C > 0$ such that for each pair of unit vectors ξ, η in H we have

$$\int_G |\langle \pi(g)\xi, \eta \rangle|^2 d\mu_{L,d}(g) \leq C$$

where $d\mu_{d,L}(g) = \frac{dg}{(1+L(g))^d}$. We say that a locally compact group has property RD with respect to L if its regular representation has property RD with respect to L .

Let $\pi : G \rightarrow U(H)$ be a unitary representation of G . For ξ and η in H , we define the matrix coefficient associated to π , as $\langle \pi(g)\xi, \eta \rangle$. Let $L^2(H)$ be the Hilbert space

Date: May 31, 2013.

1991 Mathematics Subject Classification. Primary 43; Secondary 22.

Key words and phrases. Coefficients of unitary representations, semisimple Lie groups, property RD, Furstenberg boundary.

of Hilbert-Schmidt operators. Consider the representation

$$c : G \rightarrow U(L^2(H)),$$

defined by

$$c(g)T = \pi(g)T\pi(g^{-1}),$$

for all T in $L^2(H)$. For S and T in $L^2(H)$, the matrix coefficient associated to c is $\langle c(g)S, T \rangle$. The Hilbert space $L^2(H)$ contains the Banach space of trace class operators denoted by $L^1(H)$. The restriction of the representation c to $L^1(H)$ is a Banach space representation. In Section 2 we prove the following proposition :

Proposition 1.1. *Let G be a locally compact second countable group, and let μ be a Borel measure on G which is finite on compact subsets. Let $\pi : G \rightarrow U(H)$ be a unitary representation in a Hilbert space, and let c denote the Banach space representation defined above. The following conditions are equivalent.*

- (1) *There exists $C > 0$ such that for all S, T unit vectors in $L^1(H)$ (unit for $\|\cdot\|_1$) we have*

$$\int_G |\langle c(g)S, T \rangle| d\mu(g) < C.$$
- (2) *There exists $C > 0$, such that for all unit vectors ξ, η in H we have*

$$\int_G |\langle \pi(g)\xi, \eta \rangle|^2 d\mu(g) < C.$$
- (3) *For all vectors S, T in $L^1(H)$ we have*

$$\int_G |\langle c(g)S, T \rangle| d\mu(g) < \infty.$$
- (4) *For all vectors ξ, η in H we have*

$$\int_G |\langle \pi(g)\xi, \eta \rangle|^2 d\mu(g) < \infty.$$

Take $d\mu(g) = \frac{dg}{(1+L(g))^d}$, where dg denotes a Haar measure on G , for some d in the above proposition, and notice that (1), (3) and (4) give three equivalent definitions for property RD for the unitary representation π .

Consider now a representation $\pi : G \rightarrow U(H)$ such that $H = L^2(X, m)$ where (X, m) is a measured space. We say that a function ξ is positive if $\xi \geq 0$ almost everywhere. Let

$$H^+ = \{\xi \in H \mid \xi \geq 0 \text{ almost everywhere}\},$$

be the cone of positive functions on X . In this context, we say that π is positive if for all g in G , we have $\pi(g)H^+ \subset H^+$. The following proposition proved in Section 4 brings a simple condition for property RD for positive representations:

Proposition 1.2. *Let G be a locally compact second countable group with a length function L . Let $\pi : G \rightarrow U(H)$ be a unitary representation such that $H = L^2(X, m)$ with (X, m) a measured space. Assume that π is positive. The following assertions are equivalent:*

- (1) π has property RD with respect to L i.e. there exists $d \geq 1$ and $C > 0$ such that for all ξ, η unit vectors in H , we have

$$\int_G \frac{|\langle \pi(g)\xi, \eta \rangle|^2}{(1 + L(g))^d} dg < C.$$

- (2) For all positive functions $\xi \in H$, there exists $d_\xi \geq 1$ such that

$$\int_G \frac{\langle \pi(g)\xi, \xi \rangle^2}{(1 + L(g))^{d_\xi}} dg < \infty.$$

Assume that G can be written $G = KP$, where K is a compact subgroup and P is a closed subgroup. Let $\lambda_{G/P} : G \rightarrow U(L^2(G/P))$ be the quasi-regular representation (see Subsection 3.2). The Harish-Chandra function

$$\Xi(g) = \langle \lambda_{G/P}(g)1_X, 1_X \rangle,$$

is the diagonal coefficient of $\lambda_{G/P}$ defined by the characteristic function 1_X of the space $X = G/P$. Following Gangolli and Varadarajan, [6][Definition 6.1.17], we say that a function f on the group G equipped with a length function L , verifies the *weak inequality* if there exists $C > 0$ and $d \geq 0$ such that

$$|f(g)| \leq C(1 + L(g))^d \Xi(g).$$

Theorem 1.1. *Let G be a non compact semisimple Lie group with a finite center. Then there exists a matrix coefficient of its quasi-regular representation on its Furstenberg boundary which does not verify the weak inequality .*

In Proposition 5.1 we give explicit examples of such coefficient in the special case $G = SL(2, \mathbb{R})$.

1.1. Acknowledgements. I would like to thank C. Pittet for very helpful discussions and criticisms.

2. FROM SQUARABLE REPRESENTATIONS TO INTEGRABLE REPRESENTATIONS

2.1. The spaces.

2.1.1. The Hilbert spaces. Let H be a separable complex Hilbert space with a scalar product that verifies $\langle \xi, \lambda \eta \rangle = \overline{\lambda} \langle \xi, \eta \rangle$ for all $\xi, \eta \in H$ and for all $\lambda \in \mathbb{C}$. Let $\overline{H}$ be its conjugate space. Recall $H = \overline{H}$ as an additive group. The external law for $\lambda \in \mathbb{C}$, $\xi \in H$ is defined by $\lambda \cdot \xi = \overline{\lambda} \xi$, and the scalar product is $\langle \xi, \eta \rangle_{\overline{H}} = \overline{\langle \xi, \eta \rangle}$. We consider the algebraic tensor product $\overline{H} \otimes_{alg} H$ with the scalar product defined by $\langle \xi \otimes \eta, \xi' \otimes \eta' \rangle = \langle \xi, \xi' \rangle_{\overline{H}} \langle \eta, \eta' \rangle$. We denote $\overline{H} \otimes H$ the Hilbertian completion of the algebraic tensor product.

We denote by $B(H)$ the space of bounded operators on H . Let Tr be the usual trace on $B(H)$ defined on positive bounded operators: for an orthonormal basis $(e_n)_{n \in \mathbb{N}}$ of H , for a positive operator $A \in B(H)$, we have $Tr(A) = \sum_n \langle A e_n, e_n \rangle \in [0, \infty]$. The trace Tr does not depend on the choice of the orthonormal basis. A

bounded operator T is called Hilbert-Schmidt if it verifies $Tr(|T|^2) < \infty$, where $|T| = \sqrt{T^*T}$. We denote by $L^2(H)$ the set of Hilbert-Schmidt operators. The set of Hilbert-Schmidt operators is a subset of the set of compact operators, and it is a Hilbert space under the scalar product $\langle S, T \rangle = Tr(ST^*)$. Its associated norm is $\|T\|_2 = \sqrt{Tr(TT^*)}$. It is well known that T is a Hilbert-Schmidt operator if and only if there exist two orthonormal basis $(\xi_i)_{i \in \mathbb{N}}$ and $(\eta_i)_{i \in \mathbb{N}}$ and a sequence $\alpha = (\alpha_i)_{i \in \mathbb{N}} \in l^2(\mathbb{N})$ such that the following convergent series in operator norm verifies

$$T = \sum_i \alpha_i \langle \cdot, \xi_i \rangle \eta_i.$$

Notice that if $T = \sum_i \alpha_i \langle \cdot, \xi_i \rangle \eta_i$, then $\|T\|_2 = \|\alpha\|_{l^2}$.
The map :

$$\begin{aligned} \Phi : \overline{H} \otimes H &\rightarrow L^2(H) \\ \xi \otimes \eta &\mapsto \langle \cdot, \xi \rangle \eta \end{aligned}$$

is an isomorphism of Hilbert spaces. See [11][Chap. 9.1, from § 9.1.31 to 9.1.38] and [5][Chap. 1, § 6, p. 96] for more details.

2.1.2. The Banach spaces. A bounded operator T is trace class if it verifies $Tr(|T|) < \infty$, where $|T| = \sqrt{T^*T}$. As said in the introduction, $L^1(H)$ denotes the set of trace class operators. It is a subspace of the Hilbert-Schmidt operators. It is a Banach space under the norm $\|T\|_1 = Tr(|T|)$. It is well known that T is trace class if and only if there exist two orthonormal bases $(\xi_i)_{i \in \mathbb{N}}$ and $(\eta_i)_{i \in \mathbb{N}}$ and a sequence $\alpha = (\alpha_i)_{i \in \mathbb{N}} \in l^1(\mathbb{N})$ such that the following convergent series in operator norm verifies

$$T = \sum_i \alpha_i \langle \cdot, \xi_i \rangle \eta_i.$$

Notice that if $T = \sum_i \alpha_i \langle \cdot, \xi_i \rangle \eta_i$, then $\|T\|_1 = \|\alpha\|_{l^1}$. We recall now the definition of the projective tensor product of Banach spaces. Let X (resp. Y) be a Banach space with the norm $\|\cdot\|_X$ (resp. Y). We consider the algebraic tensor product $X \otimes_{alg} Y$. The projective norm of an element u in $X \otimes_{alg} Y$ is defined by

$$p(u) := \inf \left\{ \sum_{\text{finite}} \|x_i\|_X \|y_i\|_Y, \text{ such that } u = \sum_{\text{finite}} x_i \otimes y_i \right\}.$$

The projective tensor product is defined as the completion of the algebraic tensor product for the projective norm p , and it is denoted by

$$X \widehat{\otimes} Y := \overline{X \otimes_{alg} Y}^p.$$

If H and K denote two Hilbert spaces, we observe that $H \widehat{\otimes} K \subset H \otimes K$. The map Φ restricts to an isometric Banach spaces isomorphism :

$$\begin{aligned} \Phi : \overline{H} \widehat{\otimes} H &\rightarrow L^1(H) \\ \xi \otimes \eta &\mapsto \langle \cdot, \xi \rangle \eta \end{aligned}$$

For more details see [14][Chap. 2, § 2.1, p. 12].

2.2. The representations. Let G be a locally compact group. Let $\pi : G \rightarrow U(H)$ be a unitary representation in a Hilbert space, let $\bar{\pi}$ be its conjugate unitary representation in $\bar{H}$. Let σ be the unitary representation:

$$\begin{aligned}\sigma : G &\rightarrow U(\bar{H} \otimes H) \\ g &\mapsto \bar{\pi}(g) \otimes \pi(g).\end{aligned}$$

The isomorphism Φ intertwines σ and the representation c (see Introduction):

$$c\Phi = \Phi\sigma,$$

and this equivalence restricts to an equivalence between Banach spaces representations.

Lemma 2.1. *Let U, V be a pair of unit vectors in $\bar{H} \hat{\otimes} H$. There exists a unique pair of unit vectors S, T in $L^1(H)$ such that*

$$\langle \sigma(g)U, V \rangle = \langle c(g)S, T \rangle.$$

Proof. We can write $U = \Phi(S)$ and $V = \Phi(T)$ for a unique pair $S, T \in L^1(H)$ of unit vectors because Φ is an isomorphism of Banach spaces. Furthermore, because Φ is an isomorphism of the Hilbert spaces $\bar{H} \otimes H \supset \bar{H} \hat{\otimes} H$ and $L^2(H) \supset L^1(H)$ we have

$$\begin{aligned}\langle \sigma(g)U, V \rangle &= \langle \sigma(g)\Phi(S), \Phi(T) \rangle \\ &= \langle \Phi^{-1}\sigma(g)\Phi(S), T \rangle \\ &= \langle c(g)S, T \rangle.\end{aligned}$$

□

2.3. Proof of Proposition 1.1.

2.3.1. An application of the Banach-Steinhaus theorem. The next proposition is an application of the Banach-Steinhaus theorem. It enables us to prove implication (4) $\Rightarrow$ (1) from Proposition 1.1 in the next subsection.

Proposition 2.1. *Let $B : X_1 \times X_2 \times \dots \times X_r \rightarrow \mathbb{C}$ be a multilinear map on a product of Banach spaces. If B is continuous on each variable, then B is continuous.*

Proof. By induction on r . See [13][p. 81, Corollary] for the case $r = 2$. □

If G is a locally compact second countable group, we denote by $d\mu(g)$ a Borel measure on G which is finite on compact subsets of G .

Proposition 2.2. *Let G be a locally compact second countable group. Let $\pi : G \rightarrow U(H)$ be a unitary representation in a Hilbert space and let $\sigma = \bar{\pi} \otimes \pi$. If for all $\xi, \eta \in H$ we have $\int_G |\langle \pi(g)\xi, \eta \rangle|^2 d\mu(g) < \infty$, then there exists $C > 0$ such that for all unit vectors $\xi, \xi', \eta, \eta' \in H$ we have $\int_G |\langle \sigma(g)\xi \otimes \eta, \xi' \otimes \eta' \rangle| d\mu(g) \leq C$.*

Proof. Let $G = \cup K_n$, where K_n is an exhausting sequence of compact subsets of G . Fix $\eta, \xi', \eta' \in H$, and let us define the family T_n of continuous linear operators:

$$T_n : H \rightarrow L^1(G, \mu)$$

$$\xi \mapsto (g \mapsto 1_{K_n}(g) \langle \sigma(g)\xi \otimes \eta, \xi' \otimes \eta' \rangle)$$

with 1_{K_n} the characteristic function of K_n .

We have for each $\xi \in H$,

$$\begin{aligned} \sup_n \|T_n(\xi)\|_{L^1} &= \sup_n \int_{K_n} |\langle \sigma(g)\xi \otimes \eta, \xi' \otimes \eta' \rangle| d\mu(g) \\ &= \sup_n \int_{K_n} |\langle \pi(g)\xi, \xi' \rangle| |\langle \pi(g)\eta, \eta' \rangle| d\mu(g) \\ &\leq \sup_n \left\{ \left(\int_{K_n} |\langle \pi(g)\xi, \xi' \rangle|^2 d\mu(g) \right)^{\frac{1}{2}} \left(\int_{K_n} |\langle \pi(g)\eta, \eta' \rangle|^2 d\mu(g) \right)^{\frac{1}{2}} \right\} \\ &\leq \left(\int_G |\langle \pi(g)\xi, \xi' \rangle|^2 d\mu(g) \right)^{\frac{1}{2}} \left(\int_G |\langle \pi(g)\eta, \eta' \rangle|^2 d\mu(g) \right)^{\frac{1}{2}} \\ &< \infty, \end{aligned}$$

by hypothesis. Banach Steinhaus theorem implies that $\sup_n \|T_n\| < \infty$ where $\|\cdot\|$, denotes the operator norm. Hence

$$\sup_{\{\xi, \|\xi\|=1\}} \int_G |\langle \sigma(g)\xi \otimes \eta, \xi' \otimes \eta' \rangle| d\mu(g) < \infty.$$

This proves that the multilinear form B :

$$\overline{H} \times H \times H \times \overline{H} \rightarrow \mathbb{C}$$

$$(\xi, \eta, \xi', \eta') \mapsto \int_G \langle \sigma(g)\xi \otimes \eta, \xi' \otimes \eta' \rangle d\mu(g)$$

is continuous in ξ . Analogous arguments show that B is continuous on η, ξ' and η' . Proposition 2.1 completes the proof. $\square$

2.3.2. Proof of Proposition 1.1.

Proof. (1) $\Rightarrow$ (3) and (2) $\Rightarrow$ (4) are obvious.

(1) $\Rightarrow$ (2) and (3) $\Rightarrow$ (4) are clear: take $U = \xi \otimes \xi$ and $V = \eta \otimes \eta$. Then

$$\langle \sigma(g)U, V \rangle = |\langle \pi(g)\xi, \eta \rangle|^2.$$

Lemma 2.1 and integration complete the proof.

Let us prove (4) $\Rightarrow$ (1):

Take U, V two unit vectors (for the projective norm) in $\overline{H} \hat{\otimes} H$. Write $U = \sum_k \alpha_k \xi_k \otimes \eta_k$ and $V = \sum_l \beta_l \xi'_l \otimes \eta'_l$ with $\|\alpha\|_{l^1} = 1 = \|\beta\|_{l^1}$, where $(\xi_k)_{k \in \mathbb{N}}, (\xi'_k)_{k \in \mathbb{N}}$ and $(\eta_k)_{k \in \mathbb{N}}, (\eta'_k)_{k \in \mathbb{N}}$ are orthonormal families in $\overline{H}$ and H . We have:

$$\begin{aligned} \int_G |\langle \sigma(g)U, V \rangle| d\mu(g) &= \int_G \left| \left\langle \sigma(g) \sum_k \alpha_k \xi_k \otimes \eta_k, \sum_l \beta_l \xi'_l \otimes \eta'_l \right\rangle \right| d\mu(g) \\ &\leq \sum_{k,l} |\alpha_k| |\beta_l| \int_G |\langle \sigma(g) \xi_k \otimes \eta_k, \xi'_l \otimes \eta'_l \rangle| d\mu(g). \end{aligned}$$

Thanks to Proposition 2.2, there exists $C > 0$ such that for all $k, l \in \mathbb{N}$

$$\int_G |\langle \sigma(g) \xi_k \otimes \eta_k, \xi'_l \otimes \eta'_l \rangle| d\mu(g) \leq C.$$

Hence,

$$\begin{aligned} \int_G |\langle \sigma(g)U, V \rangle| d\mu(g) &\leq \sum_{k,l} |\alpha_k| |\beta_l| C \\ &= C. \end{aligned}$$

The proof of (2) $\Rightarrow$ (3) is similar and left to the reader. $\square$

3. PROPERTY RD FOR QUASI-REGULAR REPRESENTATIONS

3.1. Property RD. We shall define property RD for a unitary representation of a locally compact group G .

Definition 3.1. A length function on G , $L : G \rightarrow \mathbb{R}_+$, is a measurable function, locally bounded (i.e. for any compact $K \subset G$, we have $\sup \{L(g), g \in K\} < \infty$) that verifies

- (1) $L(e) = 0$,
- (2) $L(g^{-1}) = L(g)$,
- (3) $L(gh) \leq L(g) + L(h)$.

Definition 3.2. A unitary representation $\pi : G \rightarrow U(H)$ of a locally compact group has property RD with respect to L if there exists $C > 0$ and $d \geq 1$ such that for all unit vectors $\xi, \eta \in H$ we have

$$\int_G \frac{|\langle \pi(g)\xi, \eta \rangle|^2}{(1 + L(g))^d} dg \leq C.$$

Remark 3.3. Assume K is a symmetric relatively compact generating set of G containing the identity. Then $L(g) = \min \{n \in \mathbb{N} : \exists k_1, \dots, k_n \text{ such that } g = k_1 \dots k_n\}$ is a length function on G . Let $C = \sup_k \{L(k), k \in K\}$. If L' is any length function on G , then $L'(g) \leq CL(g)$. Hence if π has rapid decay with respect to L' then it has rapid decay with respect to L .

3.2. Quasi-regular representations. We remind some basic facts about quasi-regular representations.

Let G be a locally compact second countable group and assume that there exists a compact subgroup K and a closed subgroup P of G such that $G = KP$. We denote by Δ_P the right-modular function of P . We extend to G the map Δ_P of P as $\Delta : G \rightarrow \mathbb{R}_+^*$ with $\Delta(g) = \Delta(kp) := \Delta_P(p)$. It is well defined because $K \cap P$ is compact (observe that $\Delta_P|_{K \cap P} = 1$). Notice that for all $g \in G$, the function $x \in G/P \mapsto \frac{\Delta(gx)}{\Delta(x)} \in \mathbb{R}_+^*$ is well defined. The quotient G/P carries a unique quasi-invariant measure ν , such that the Radon-Nikodym derivative at $(g, x) \in G \times G/P$ denoted by $\kappa(g, x) = \frac{dg_*\nu}{d\nu}(x)$ with $g_*(A) = \nu(g^{-1}A)$, satisfies $\frac{dg_*\nu}{d\nu}(x) = \frac{\Delta(gx)}{\Delta(x)}$ for all $g \in G$ and $x \in G/P$. Consider the quasi-regular representation $\lambda_{G/P} : G \rightarrow U(L^2(G/P))$ associated to P , defined by $(\lambda_{G/P}(g)\xi)(x) = \kappa(g^{-1}, x)^{\frac{1}{2}}\xi(g^{-1} \cdot x)$. We denote by X the space G/P , and by 1_X the characteristic function of G/P . We refer to [1][Appendix B, Lemma B.1.3, p. 344-345] for more details.

Definition 3.4. *The Harish-Chandra function $\Xi : G \rightarrow (0, \infty)$ is defined as*

$$\Xi(g) := \langle \lambda_{G/P}(g)1_X, 1_X \rangle.$$

As $\lambda_{G/P}$ is a unitary representation, we have $\Xi(g) = \Xi(g^{-1})$ for all $g \in G$. The Harish-Chandra function is a stable matrix coefficient in the following sense. Let Γ be a discrete subgroup of a locally compact group G . Let $\pi : G \rightarrow U(H)$ be a unitary representation. A matrix coefficient, $\langle \pi(g)\xi, \eta \rangle$ is *stable* on G relatively to Γ if there exists a relatively compact neighborhood V of the identity $e \in G$ and $C, c > 0$ such that

$$c |\langle \pi(\gamma)\xi, \eta \rangle| \leq |\langle \pi(\gamma g)\xi, \eta \rangle| \leq C |\langle \pi(\gamma)\xi, \eta \rangle|$$

for all $g \in V$ and for all $\gamma \in \Gamma$.

Lemma 3.5. ([6, Proposition 4.6.3, p. 159].) *The Harish-Chandra function is stable relatively to any discrete subgroup.*

Proof. Let V be a relatively compact neighborhood of e in G , sufficiently small such that $\gamma \cdot V \cap \gamma' \cdot V = \emptyset$ for all $\gamma \neq \gamma' \in \Gamma$ a discrete subgroup of G . By definition of the Harish-Chandra function, we have $\Xi(\gamma g) = \langle \lambda_{G/P}(\gamma g)1_X, 1_X \rangle = \langle \lambda_{G/P}(g)1_X, \lambda_{G/P}(\gamma^{-1})1_X \rangle$. That is $\Xi(\gamma g) = \int_X \kappa(g^{-1}, x)^{\frac{1}{2}} \kappa(\gamma, x)^{\frac{1}{2}} d\nu(x)$. The function $(g, x) \in G \times G/P \mapsto \kappa(g, x)^{\frac{1}{2}}$ is a strictly positive continuous function. Therefore, as $\bar{V}$ and G/P are compact, there exist $C, c > 0$ such that for all $g \in \bar{V}$ and for all $x \in G/P$, we have $c \leq \kappa(g^{-1}, x)^{\frac{1}{2}} \leq C$. Hence, for all $\gamma \in \Gamma$, for all $g \in V$,

$$c\Xi(\gamma) \leq \Xi(\gamma g) \leq C\Xi(\gamma).$$

□

Lemma 3.6. *Let Γ be a discrete subgroup of a locally compact group G . Let $\pi : G \rightarrow U(H)$ be a unitary representation, and let L be a length function on G . If $\langle \pi(g)\xi, \eta \rangle$ is a stable coefficient relatively to Γ , then there exists a constant $C \geq 1$ such that*

$$\sum_{\gamma \in \Gamma} \frac{|\langle \pi(\gamma)\xi, \eta \rangle|^2}{(1 + L(\gamma))^d} \leq C \int_G \frac{|\langle \pi(g)\xi, \eta \rangle|^2}{(1 + L(g))^d} dg.$$

Proof. Let V be a relatively compact neighbourhood of the neutral element of G , such that $\gamma \cdot V \cap \gamma' \cdot V = \emptyset$ for all $\gamma, \gamma' \in \Gamma$ such that $\gamma \neq \gamma'$. Take a matrix coefficient $\langle \pi(\gamma)\xi, \eta \rangle$ which is stable. We have

$$\begin{aligned} \int_G \frac{|\langle \pi(g)\xi, \eta \rangle|^2}{(1 + L(g))^d} dg &\geq \sum_{\gamma} \int_{\gamma \cdot V} \frac{|\langle \pi(g)\xi, \eta \rangle|^2}{(1 + L(g))^d} dg \\ &= \sum_{\gamma} \int_V \frac{|\langle \pi(\gamma x)\xi, \eta \rangle|^2}{(1 + L(\gamma x))^d} dx \\ &\geq \sum_{\gamma} \int_V c^2 \frac{|\langle \pi(\gamma)\xi, \eta \rangle|^2}{(1 + L(\gamma x))^d} dx \\ &\geq \epsilon \sum_{\gamma} \frac{|\langle \pi(\gamma)\xi, \eta \rangle|^2}{(1 + L(\gamma))^d}, \end{aligned}$$

for some positive constant ϵ . □

Combining Lemmas 3.5 and 3.6, we obtain the following

Proposition 3.1. *Let G be a locally compact group decomposed as $G = KP$ where K is a compact subgroup and P is a closed subgroup. Let Γ be a discrete subgroup of G and let L be a length function on G . Let $d \geq 1$. There exists a constant $C > 1$ such that*

$$\sum_{\gamma \in \Gamma} \frac{\Xi^2(\gamma)}{(1 + L(\gamma))^d} \leq C \int_G \frac{\Xi^2(g)}{(1 + L(g))^d} dg.$$

3.3. Linearization. The representation $\sigma : G \rightarrow U(\overline{H} \otimes H)$ introduced in Subsection 2.2 satisfies for all $\xi, \eta \in H$:

$$\langle \sigma(g)\xi \otimes \xi, \eta \otimes \eta \rangle = |\langle \pi(g)\xi, \eta \rangle|^2.$$

The representation σ can be used to give a short and elementary proof of the following known result, due to C. Herz.

Theorem 3.7. ([8], C. Herz.) *Let G be a connected semisimple Lie group with finite center. Then G has property RD with respect to the length function associated to any left-invariant Riemannian metric on G .*

See [8], [4], [2] for proofs.

4. EQUIVALENT DEFINITIONS FOR PROPERTY RD AND THE WEAK INEQUALITY

4.1. Positivity.

Lemma 4.1. *Let G be a locally compact second countable group with a length function L . Let $\pi : G \rightarrow U(H)$ be a unitary representation with $H = L^2(X, m)$ where (X, m) is a measured space. Assume that π is positive (π preserves the cone of positive functions). The following assertions are equivalent.*

(1) *There exists $d \geq 1$ such that for all vectors $\xi, \eta \in H$ we have*

$$\int_G \frac{|\langle \pi(g)\xi, \eta \rangle|^2}{(1 + L(g))^d} dg < \infty.$$

(2) *There exists $d \geq 1$ such that for all positive vectors $\xi \in H$ we have*

$$\int_G \frac{\langle \pi(g)\xi, \xi \rangle^2}{(1 + L(g))^d} dg < \infty.$$

Proof. (1) $\Rightarrow$ (2) is obvious.

Let us prove (2) $\Rightarrow$ (1). Observe first that by decomposition of a real valued function ξ into its positive and negative part we have that $|\xi_+ - \xi_-| \leq \xi_+ + \xi_-$. By positivity of π , for all $g \in G$ we have $\pi(g)|\xi_+ - \xi_-| \leq \pi(g)\xi_+ + \pi(g)\xi_-$. Using the decomposition of a complex valued function into its real and imaginary parts, and the decomposition of a real valued function into its positive and negative part, we obtain, according to the above observation, that $|\langle \pi(g)\xi, \eta \rangle|$ is less than or equal to a linear combination of matrix coefficients $\langle \pi(g)\xi', \eta' \rangle$ with ξ', η' positive vectors in H . Now observe that for positive vectors,

$$\langle \pi(g)\xi', \eta' \rangle \leq \langle \pi(g)(\xi' + \eta'), \xi' + \eta' \rangle.$$

Integration and Cauchy-Schwarz inequality complete the proof. $\square$

Here is the proof of Proposition 1.2:

Proof. (1) $\Rightarrow$ (2) is clear.

(2) $\Rightarrow$ (1). Thanks to the equivalence between (2) and (4) in Proposition 1.1 with $d\mu(g) = \frac{dg}{(1+L(g))^d}$ for some $d \geq 1$, it is enough to prove that there exists $d \geq 1$, such that for each pair of vectors $\xi, \eta \in H$ we have $\int_G \frac{|\langle \pi(g)\xi, \eta \rangle|^2}{(1+L(g))^d} dg < \infty$. According to Lemma 4.1 it is equivalent to prove that there exists $d \geq 1$ such that for any positive function $\xi \in L^2(X, m)$ we have $\int_G \frac{\langle \pi(g)\xi, \xi \rangle^2}{(1+L(g))^d} dg < \infty$. We give a proof by contraposition.

Assume that for each $n \in \mathbb{N}$, there exists a unit positive vector ξ_n such that

$$\int_G \frac{\langle \pi(g)\xi_n, \xi_n \rangle^2}{(1 + L(g))^n} dg = \infty.$$

Take a sequence $(a_n)_{n \in \mathbb{N}}$ of strictly positive real numbers such that the series $\sum_n a_n$ converges. We consider

$$\xi = \sum_n a_n \xi_n$$

which is a well defined element of H . We can assume that $\xi \neq 0$. We can replace ξ by $\frac{\xi}{\|\xi\|}$ which is a unit vector, so we assume that ξ is a unit vector. Let d be a positive real number. Let n be an integer such that $n \geq d$. Notice that $\langle \pi(g)\xi_n, \xi_m \rangle \geq 0$ for all $n, m \in \mathbb{N}$ and for all g in G . Hence,

$$\begin{aligned} \int_G \frac{\langle \pi(g)\xi, \xi \rangle^2}{(1 + L(g))^d} dg &\geq \int_G \frac{\langle \pi(g)\xi, \xi \rangle^2}{(1 + L(g))^n} dg \\ &\geq a_n^4 \int_G \frac{\langle \pi(g)\xi_n, \xi_n \rangle^2}{(1 + L(g))^n} dg = \infty. \end{aligned}$$

Finally we have found a unit vector ξ , such that for all $d \geq 1$ we have

$$\int_G \frac{\langle \pi(g)\xi, \xi \rangle^2}{(1 + L(g))^d} dg = \infty$$

as required. $\square$

4.2. The weak inequality.

Definition 4.2. Let G be a locally compact group with a length function L , which can be written $G = KP$ where K is a compact subgroup and P is a closed subgroup. A function on G is said to satisfy the weak inequality if there exists $C > 0$ and $d \geq 0$ such that for all $g \in G$

$$|f(g)| \leq C(1 + L(g))^d \Xi(g).$$

For more details about the weak inequality we refer to [6][§ 6.1.17, p. 258] and [16][§ 8.4, p. 231-232]

We prove Theorem 1.1:

Proof. Let $u \in G$ be a parabolic element. Consider $\Gamma \cong \mathbb{Z}$ the cyclic subgroup generated by u . Observe that Γ is discrete and is a group of exponential growth with respect to the restriction of L to Γ . But an amenable group satisfies property RD if and only if it is a group of polynomial growth, see [9][Corollary 3.1.8]. Hence, according to Shalom's Lemma, [15][Lemma 2.3], the restriction of $\lambda_{G/P}$ to Γ does not satisfy property RD. By Proposition 1.2 applied to Γ , we can find a positive vector $\xi \in L^2(G/P)$ such that for all $d > 1$ we have

$$\sum_{\gamma \in \Gamma} \frac{\langle \lambda_{G/P}(\gamma)\xi, \xi \rangle^2}{(1 + L(\gamma))^d} = \infty.$$

We claim that the weak inequality fails for the coefficient $\langle \lambda_{G/P}(g)\xi, \xi \rangle$. Assume on the contrary that it holds. There would exist C_ξ and d_ξ such that for all $g \in G$ $\langle \lambda_{G/P}(g)\xi, \xi \rangle \leq C_\xi(1 + L(g))^{d_\xi}\Xi(g)$. For any $d_0 > 0$, we would have:

$$\begin{aligned} \sum_{\gamma \in \Gamma} \frac{\langle \lambda_{G/P}(\gamma)\xi, \xi \rangle^2}{(1 + L(\gamma))^{2d_\xi + d_0}} &\leq C_\xi^2 \sum_{\gamma \in \Gamma} \frac{(1 + L(\gamma))^{2d_\xi}\Xi(\gamma)^2}{(1 + L(\gamma))^{2d_\xi + d_0}} \\ &= C_\xi^2 \sum_{\gamma \in \Gamma} \frac{\Xi(\gamma)^2}{(1 + L(\gamma))^{d_0}} \\ &\leq C_\xi^2 C \int_G \frac{\Xi(g)^2}{(1 + L(g))^{d_0}} dg, \end{aligned}$$

where the last inequality follows from Proposition 3.1.

According to Theorem 3.7, there exists $d_0 > 0$ such that $\int_G \frac{\Xi(g)^2}{(1 + L(g))^{d_0}} < \infty$. It would follow

$$\sum_{\gamma \in \Gamma} \frac{\langle \lambda_{G/P}(\gamma)\xi, \xi \rangle^2}{(1 + L(\gamma))^{2d_\xi + d_0}} < \infty.$$

This is a contradiction. □

4.3. Examples of coefficients that satisfy the weak inequality.

4.3.1. *A trivial example:* $L^\infty(G/P)$. If $\xi, \eta \in L^\infty(G/P)$, we have

$$|\langle \lambda_{G/P}(g)\xi, \eta \rangle| \leq \|\xi\|_\infty \|\eta\|_\infty \Xi(g).$$

4.3.2. *The K -finite vectors.* Let G be a locally compact second countable unimodular group, such that $G = KP$ with K a compact subgroup of G , and P a closed subgroup of G . Let $\pi : G \rightarrow U(H)$ be a unitary representation of G . We say that ξ in H is a K -finite vector if

$$\dim \text{Span} \{ \pi(k)\xi, k \in K \} < \infty.$$

We denote by $d(\xi)$ the dimension of $\text{Span} \{ \pi(k)\xi, k \in K \}$. Consider $\lambda_{G/P}$ the quasi-regular representation associated to P . Then for ξ, η two K -finite vectors we have

$$|\langle \lambda_{G/P}(g)\xi, \eta \rangle| \leq \|\xi\| \|\eta\| d(\xi)^{\frac{1}{2}} d(\eta)^{\frac{1}{2}} \Xi(g).$$

For a more general statement we refer to [15][Proposition 2.5].

5. A NON-EXAMPLE IN $SL_2(\mathbb{R})$

Let $G = SL_2(\mathbb{R})$. We consider the Iwasawa decomposition $G = KAN$, where

$$\begin{aligned} K = SO_2(\mathbb{R}) &= \left\{ R_\theta = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix}, \theta \in]-\pi, \pi] \right\}, \\ A &= \left\{ \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \lambda \in \mathbb{R}_+^* \right\} \end{aligned}$$

and

$$N = \left\{ \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, t \in \mathbb{R} \right\}.$$

Let $M = C_K(A)$ be the centralizer of A in K . We have $M = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \right\}$. Notice that M is central in G and A normalizes N . Let P be the group $P = MAN$. Hence $P = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix}, a \in \mathbb{R}^*, b \in \mathbb{R} \right\}$. We have $G = KP$.

We identify $G/P \cong \mathbb{S}^1/\{\pm 1\}$. The Lebesgue measure, $\frac{d\theta}{\pi}$ on $\mathbb{S}^1/\{\pm 1\}$ is quasi-invariant under the action of G , and is invariant under the action of K , i.e. by rotation. We identify the class of functions $\xi \in L^2(G/P)$ with the class of π -periodic functions $\xi : \mathbb{R} \rightarrow \mathbb{C}$, that verifies

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} |\xi(\theta)|^2 \frac{d\theta}{\pi} < \infty.$$

An element $g \in G$ can be written $g = kp$, with $k \in SO_2(\mathbb{R})$, and where $p = \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix}$ with $a \in \mathbb{R}^*$, and $b \in \mathbb{R}$. If Δ_P denotes the modular function on P , we have

$$\Delta(kp) := \Delta_P(p) = \Delta_P \left(\begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \right) = a^{-2}.$$

Consider the hyperbolic distance d , on the hyperbolic plane

$$\mathbb{H} = \{z \in \mathbb{C}, \text{Im}(z) > 0\}.$$

The formula $L(g) = d(g \cdot i, i)$ defines a length function on G where the action $G \curvearrowright \mathbb{H}$ is $g \cdot z = \frac{az+b}{cz+d}$ with $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R})$ and $z \in \mathbb{H}$.

The quasi-regular representation associated to P is given for all $\xi \in L^2(G/P)$, for all $g \in G$, for almost everywhere $\theta \in G/P$, by:

$$\lambda_{G/P}(g)\xi(\theta) = \Delta(g^{-1}R_\theta)^{\frac{1}{2}}\xi(g^{-1} \cdot \theta).$$

Define for $s \geq 0$ the Sobolev space

$$H^s(\mathbb{S}^1) := \left\{ \xi \in L^2(\mathbb{S}^1) \mid \sum_n |c_n|^2 (1+n^2)^s < \infty \right\},$$

where the sequence $(c_n)_{n \in \mathbb{Z}}$ is the sequence of the Fourier coefficients of ξ on $\mathbb{S}^1$. The space $H^s(\mathbb{S}^1)$ is a Hilbert space for the scalar product

$$\langle \xi, \eta \rangle = \left(\sum_{n \in \mathbb{Z}} (1+n^2)^s \overline{c_n} d_n \right)$$

where the $(c_n)_{n \in \mathbb{Z}}$ and the $(d_n)_{n \in \mathbb{Z}}$ are respectively the Fourier coefficients of ξ and η . Notice that for $s > \frac{1}{2}$, we have $H^s(\mathbb{S}^1) \subset C(\mathbb{S}^1)$. Let δ such that $\frac{1}{4} < \delta < \frac{1}{2}$. Define

$$\xi_\delta : \theta \in [-\frac{\pi}{2}, \frac{\pi}{2}] \mapsto \frac{1}{\theta^\delta} \in L^2(G/P).$$

The following proposition can be regarded as a more precise statement, in the case of $SL(2, \mathbb{R})$, of Theorem 1.1.

Proposition 5.1. *Let $0 \leq s < \frac{1}{4}$. Let $\frac{1}{4} < \delta < \frac{1}{2} - s$. Then $\xi_\delta \in H^s(\mathbb{S}^1)$ and the coefficient $\langle \lambda_{G/P}(g)\xi_\delta, \xi_\delta \rangle$ does not satisfy the weak inequality.*

See [3] for the proof.

Remark 5.1. *To sum up, if $0 \leq s < \frac{1}{4}$ the weak inequality fails, if $s \geq \frac{1}{2}$ the weak inequality is obviously true for all coefficients defined by vectors of $H^s(\mathbb{S}^1)$, because $H^s(\mathbb{S}^1) \subset C(\mathbb{S}^1)$, and if $\frac{1}{4} \leq s \leq \frac{1}{2}$ we don't know.*

REFERENCES

- [1] B. Bekka, P. de la Harpe, and A. Valette, *Kazhdan's property (T)*, New Mathematical Monographs, vol. 11, Cambridge University Press, Cambridge, 2008.
- [2] A. Boyer, *Semisimple Lie groups satisfy property RD, a short proof*, submitted, 2013.
- [3] A. Boyer, *PhD thesis*, in preparation, 2013.
- [4] I. Chatterji, C. Pittet, and L. Saloff-Coste, *Connected Lie groups and property RD*, Duke Math. J. 137 (2007), no. 3, 511-536.
- [5] J. Dixmier, *Les algèbres d'opérateurs dans l'espace hilbertien*, Gauthiers-Villars, Paris, 1964.
- [6] R. Gangolli, V.S. Varadarajan, *Harmonic Analysis of Spherical Functions on Real Reductive Groups*, Springer-Verlag, New-York, 1988.
- [7] U. Haagerup, *An example of a nonnuclear C^* -algebra which has the metric approximation property*, Invent. Math. 50 (1978/79), no. 3, 279-293.
- [8] C. Herz, *Sur le phénomène de Kunze-Stein*, C. R. Acad. Sci. Paris Sr. A-B 271 (1970), A491-A493.
- [9] P. Jolissaint, *Rapidly decreasing functions in reduced C^* -algebras of groups*, Trans. Amer. Math. Soc. 317 (1990), no. 1, 167-196.
- [10] V. Lafforgue, *K-théorie bivariante pour les algèbres de Banach et conjecture de Baum-Connes*, Invent. Math. 149 (2002), no. 1, 1-95 (French).
- [11] T-W. Palmer, *Banach Algebras and the General Theory of $*$ -Algebras: Volume 2*, Encyclopedia of Mathematics and its Applications, vol. 49, Cambridge University Press, Cambridge, (2001)
- [12] M. Perrone, *Rapid decay and weak containment of unitary representations*, unpublished notes, (2009)
- [13] M. Reed, B. Simon, *I: Functional Analysis, Volume 1 (Methods of Modern Mathematical Physics)*, Academic press, (1980)
- [14] R-A. Ryan, *Introduction to tensor product of Banach spaces*, Springer Monographs in Mathematics, (2002)
- [15] Y. Shalom, *Rigidity, unitary representations of semisimple groups, and fundamental groups of manifolds with rank one transformation group*, Ann. of Math. (2) 152 (2000), no. 1, 113-182.

- [16] V. S. Varadarajan, *Reflections on quanta, symmetries, and supersymmetries*, Springer, New York, (2011)

LATP, CMI, AIX-MARSEILLE UNIV. FRANCE

E-mail address: `aboyer@cmi.univ-mrs.fr`